

MATHEMATICS (FINAL)

1. The system of equations
$$\begin{pmatrix} 1 & 3 & -2 \\ -3 & 0 & -5 \\ 2 & 5 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}$$

- A. has no solution
- B. has one and only one solution
- C. has infinite number of solutions
- D. None of the above

2. If a and b are real numbers, then $\sup\{a, b\} =$

A. $\frac{a+b-|a-b|}{2}$

B. $\frac{a-b-|a-b|}{2}$

C. $\frac{a-b+|a+b|}{2}$

D. $\frac{a+b+|a-b|}{2}$

3. If, for $x \in \mathbb{R}$, $\varphi(x)$ denotes the integer closest to x (if there are two such integers take the larger one), then $\int_{10}^{12} \varphi(x) dx$ equals

- A. 22
- B. 11
- C. 20
- D. 12

4. The eigen values of the matrix $\begin{pmatrix} 2 & 1 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 2 \end{pmatrix}$ are

A. 1,1,1

B. 2,2,2

C. -2, -2, -2

D. -1, -1, -1

5. The value of $\sum_{n=1}^{\infty} \frac{(-1)^{(n+1)}}{n}$ is

A. $\log 2$

B. e^2

C. 0

D. 1

6. The rank of the matrix $\begin{pmatrix} 0 & 0 & 0 & 0 & -2 \\ 1 & 0 & 0 & 0 & -4 \\ 0 & 1 & 0 & 0 & 5 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 13 \end{pmatrix}$ is

A. 5

B. 4

C. 3

D. 2

7. $\sum_{n=1}^{\infty} \frac{2^n}{(n-1)!}$ is

A. e

B. $2e$

C. $2e^2$

D. $2e^{-2}$

8. If $A = \begin{bmatrix} \alpha & 2 \\ 2 & \alpha \end{bmatrix}$, and $|A^2| = 125$, then the value of α is
- A. ± 5
 - B. ± 3
 - C. ± 2
 - D. ± 1
9. The algebraic equation $x^n + y^n = z^n$ has no integer solution when
- A. $n > 2$
 - B. $n = 2$
 - C. $n = 0$
 - D. $n < 2$
10. Which of the function $f : R \rightarrow R$ is one-one and onto?
- A. $f(x) = x^3 + 2$
 - B. $f(x) = \sin x$
 - C. $f(x) = \cos x$
 - D. $f(x) = x^4 - x^2$
11. Let $P(x)$ be a non-constant polynomial such that $P(n) = P(-n)$ for all $n \in N$. Then $P'(0)$ is
- A. -1
 - B. 1
 - C. 0
 - D. -2

12. If $A = \begin{bmatrix} 3 & 1 \\ -1 & 2 \end{bmatrix}$, then $A^2 - 5A + 7I$ is equal to
- A. $\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$
- B. $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
- C. $2 \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$
- D. None of the above
13. If A is any non-identity 3×3 matrix such that $A^2 = A$, then A is
- A. Singular
- B. Non singular
- C. Regular
- D. None of the above
14. If every cross-section of a bounded surface in three dimensions is a circle, then the surface must be a
- A. cylinder
- B. sphere
- C. cone
- D. third-degree surface
15. For any complex number z , the minimum value of $|z| + |z - 1|$ is
- A. 1
- B. 0
- C. $1/2$
- D. $3/2$

16. The number having a recurring decimal representation $1.414141\dots$ is
- A. real but irrational
 - B. not real
 - C. rational
 - D. neither rational nor real
17. The derivative with respect to x of the product $(1+x)(1+x^2)(1+x^4)(1+x^8)+\dots+(1+x^{56})$ at $x=0$ is
- A. 0
 - B. 1
 - C. 8
 - D. 56
18. If α , β and γ are roots of the equation $x^3 + px^2 + qx + r = 0$, then $\alpha^2 + \beta^2 + \gamma^2$ is equal to
- A. $p^2 - 2q$
 - B. $p^2 + 2q$
 - C. $2p + q^2$
 - D. $2p - q^2$
19. Given that $2+i\sqrt{3}$ is one of $x^3 - 5x^2 + 11x - 7 = 0$ then the other roots are
- A. $2-i\sqrt{3}, -1$
 - B. $2-i\sqrt{3}, 1$
 - C. $2+i\sqrt{3}, 1$
 - D. None of the above

20. The quadratic function on a single variable attains its maximum value 5 at 3. The function is
- A. $ax^2 - 6ax + 9a + 5, a < 0$
 - B. $ax^2 - 6ax + 9a + 5, a > 0$
 - C. $ax^2 + 6ax + 9a + 5, a < 0$
 - D. None of the above
21. Let n be a two digit number. $P(n)$ is the product of the digits of n and $S(n)$ is the sum of the digits of n . If $n = P(n) + S(n)$ then the unit digit of n is
- A. 1
 - B. 5
 - C. 7
 - D. 9
22. The value of $\sqrt{20 + \sqrt{20 + \sqrt{20 + \dots}}}$ is
- A. -4
 - B. 5
 - C. 4
 - D. -5
23. If A, B, C are three sets with cardinality m, p, q respectively such that $B \cap C = \phi$, then the cardinality of $(A \times B) \cup (A \times C)$ equals
- A. mpq
 - B. $m(p+q)$
 - C. $m(p+q-pq)$
 - D. $m+pq$

24. Which among the following is not a group under usual multiplication?
- A. $i - \{0\}$
 - B. $\mathbb{R} - \{0\}$
 - C. $\mathbb{R}^+$
 - D. $\mathbb{Y}$
25. If $\log_{27} x = \log_3 27$, then x is
- A. 27
 - B. 3
 - C. 3^{27}
 - D. 27^3
26. If ϕ is a homomorphism of a group G onto a group $\overline{G}$ with kernel K , then G/K is isomorphic to
- A. G
 - B. $\overline{G}$
 - C. $\overline{G}/\overline{K}$
 - D. None of the above
27. If G is a group and for $a \in G$ no positive integer m exists such that $a^m = e$, then the order of a is
- A. finite
 - B. m
 - C. $m + 1$
 - D. infinite

28. The derivative of e^t with respect to $\sqrt{t}$ is

A. $\frac{e^t}{2\sqrt{t}}$

B. $\frac{2\sqrt{t}}{e^t}$

C. $2\sqrt{t}e^t$

D. $2\sqrt{te^t}$

29. $\int_0^1 \int_0^1 \frac{1}{\sqrt{(1-x^2)} \sqrt{(1-y^2)}} dx dy$, is equal to

A. $\frac{\pi^2}{2}$

B. $\frac{\pi^2}{3}$

C. $\frac{\pi^2}{4}$

D. None of the above

30. $\int_0^{\frac{\pi}{2}} \sin^2 x \cos^2 x dx$ is equal to

A. $\frac{\pi}{8}$

B. $\frac{\pi}{16}$

C. $\frac{\pi}{32}$

D. 1

31. $\int_0^{\infty} \frac{1}{(1+x^2)^4} dx$ is equal to

A. $\frac{-5\pi}{32}$

B. $\frac{5\pi}{16}$

C. $\frac{5\pi}{32}$

D. $\frac{-5\pi}{16}$

32. $\int_0^{\frac{\pi}{6}} \cos^2 \frac{\pi}{2}$ is equal to

A. $\frac{\pi}{12} - \frac{1}{4}$

B. $\frac{\pi}{12} + \frac{1}{4}$

C. $\frac{1}{4} - \frac{\pi}{2}$

D. $\frac{1}{4} + \frac{\pi}{2}$

33. The series $\sum \frac{1}{n(\log n)^p}$ is divergent if

A. $p > 1$

B. $p \leq 1$

C. $p < 1$

D. $p = 1$

34. A non-decreasing sequence which is bounded above is

- A. divergent
- B. convergent
- C. oscillating
- D. unbounded

35. Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be defined by

$$f(x) = \begin{cases} \frac{1 - \cos 4x}{x^2} & \text{if } x < 0 \\ a, & \text{if } x = 0 \\ \frac{\sqrt{x}}{\sqrt{16 + \sqrt{x}} - 4} & \text{if } x > 0 \end{cases}$$

If f is continuous for all x , then the value of a is

- A. 0
- B. 4
- C. 8
- D. 5

36. The cardinal number of the empty set is

- A. 1
- B. 0
- C. ∞
- D. -1

37. Every closed and bounded set in $\mathbb{R}^n$ is

- A. empty
- B. open
- C. compact
- D. convex

38. In which subspace the sequence $\left\{\frac{1}{n}\right\}$ is Cauchy but not convergent
- A. $[0,1]$
 - B. $[0,1)$
 - C. $(0,1]$
 - D. $(0,1)$
39. The collection of open intervals $\left(\frac{1}{n}, \frac{2}{n}\right), n = 1, 2, 3, \dots$ is an open covering of the interval
- A. $(0,1)$
 - B. $(1,2)$
 - C. $(0,\infty)$
 - D. $(1,\infty)$
40. The set of all rational numbers is a
- A. empty set
 - B. finite set
 - C. countable set
 - D. uncountable set
41. $\int_0^{2\pi} \frac{dx}{(2 + \cos x)}$ is equal to
- A. $\frac{2\pi}{\sqrt{3}}$
 - B. $\frac{\pi}{\sqrt{3}}$
 - C. $\frac{2\pi}{-3}$

D. None of the above

42. The whole length of the asteroid $x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}$ is

A. $6a$

B. $8a$

C. $4a$

D. None of the above

43. The condition for the point (x, y) to lie on the straight line joining the points $(0, b)$ and $(a, 0)$ is

A. $\frac{x}{a} + \frac{y}{b} = 1$

B. $\frac{x}{a} - \frac{y}{b} = 1$

C. $\frac{x}{a^2} + \frac{y}{b^2} = 1$

D. None of the above

44. The centroid of the triangle whose vertices are $(2, 4, -3)$, $(-3, 3, -5)$ and $(-5, 2, -1)$ is

A. $(-2, -3, -3)$

B. $(-3, 3, -2)$

C. $(3, -2, -3)$

D. $(-2, 3, -3)$

45. The triangle whose vertices are $(-1, 1, 0)$, $(3, 2, 1)$ and $(1, 3, 2)$ is

A. isosceles triangle

B. right-angle triangle

C. equilateral triangle

D. None of the above

46. The coordinates of the point at which the line joining the points $(4, 3, 1)$ and $(1, -2, 6)$ meets the plane $3x - 2y - z + 3 = 0$ is
- A. $(-2, -7, 11)$
 - B. $(-2, 7, 11)$
 - C. $(-2, -7, -11)$
 - D. $(2, 7, 11)$
47. The center of the sphere $x^2 + y^2 + z^2 - 6x + 8y - 10z + 1 = 0$ is
- A. $(5, -4, 3)$
 - B. $(3, 4, -5)$
 - C. $(-5, -4, -3)$
 - D. $(3, -4, 5)$
48. The equation of the right circular cone with its vertex at the origin, axis along z-axis and semi-vertical angle α is
- A. $x^2 + y^2 = z^2 \tan^2 \alpha$
 - B. $x^2 - y^2 = z^2 \tan^2 \alpha$
 - C. $x^2 + y^2 = z \tan^2 \alpha$
 - D. $x^2 - y^2 = z \tan^2 \alpha$
49. $\nabla \times (\nabla \times A)$ is equal to
- A. 0
 - B. $-\nabla^2 A + \nabla(\nabla \cdot A)$
 - C. $\nabla^2 A + \nabla(\nabla \cdot A)$
 - D. $(\nabla \times \nabla) \times A$

50. The curl of $\vec{F} = x^2\vec{i} + y^2\vec{j} + z^2\vec{k}$ at $(1, 2, -3)$ is
- A. $\vec{i} + 2\vec{j} - 3\vec{k}$
 - B. $\vec{i} - 2\vec{j} + 3\vec{k}$
 - C. $2\vec{i} - 4\vec{j} + 6\vec{k}$
 - D. $2\vec{i} + 4\vec{j} - 6\vec{k}$
51. Let $F = \frac{-yi + xj}{x^2 + y^2}$. Then $\nabla \times F$ is
- A. 0
 - B. 1
 - C. -1
 - D. None of the above
52. The probability mass function or probability density function for which the mean in units and the variance in square units are same is
- A. Binomial
 - B. Poisson
 - C. Standard Normal
 - D. Geometric
53. The chance of getting at least 9, in a single throw with two dice is
- A. $\frac{4}{36}$
 - B. $\frac{3}{36}$
 - C. $\frac{5}{18}$
 - D. $\frac{1}{18}$

54. A random variable X has a probability density function $f(x) = \frac{C}{1+x^2}$, $-\infty < x < \infty$. Then the value of C is
- A. π
 - B. 1
 - C. $\frac{1}{\pi}$
 - D. $\frac{2}{\pi}$
55. If $f(z)$ and $f(\bar{z})$ are analytic in a region D , then
- A. $f(z)$ is constant in D
 - B. $f(z)$ is continuous in D
 - C. $f(z)$ is not differentiable everywhere in D
 - D. $f(z) = |z|$
56. The value of $\int_C \frac{z^2+1}{z^2-1} dz$, where C is a circle of unit radius with center at $z=1$ is
- A. 0
 - B. $2\pi i$
 - C. $-2\pi i$
 - D. 1

57. If $(x + iy)^{\frac{1}{3}} = a + ib$, then the value of $\frac{x}{a} + \frac{y}{b}$ is
- A. $4(a^2 - b^2)$
B. $4ab$
C. $4(a^2 + b^2)$
D. $5ab$
58. The value of $(1 + i)(1 + i^2)(1 + i^3)(1 + i^4) \dots (1 + i^{50})$ is
- A. 50
B. 1
C. 0
D. i
59. If $\omega = \frac{z}{z - i}$ and $|\omega| = 1$, then z lies on
- A. a circle
B. an ellipse
C. a parabola
D. a straight line
60. The residue of $\frac{z^2}{(z - 1)(z - 2)(z - 3)}$ at $z = 1$ is
- A. -8
B. $1/2$
C. -6
D. 0

61. If $u(x, y) = xy$ is a harmonic function, a harmonic conjugate of u is

A. $\frac{x^2}{2}$

B. $x^2 + y^2$

C. $\frac{x^2}{2} + \frac{y^2}{2}$

D. $\frac{y^2}{2} - \frac{x^2}{2}$

62. The solution of a homogeneous initial value problem with constant coefficient is $y = 3xe^{2x} + 6\cos 4x$. Then the least possible order of the differential equation is

A. 4

B. 5

C. 6

D. 3

63. The differential equation of the curve $y = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$ is

A. $\frac{dy}{dx} = x$

B. $\frac{dy}{dx} = -x$

C. $\frac{dy}{dx} = y$

D. $\frac{dx}{dy} = y$

64. The differential equation of all circles which pass through the origin and whose centers are on the x-axis is
- A. $y^2 + x^2 - 2xy \frac{dy}{dx} = 0$
- B. $x^2 - y^2 + 2x \frac{dy}{dx} = 0$
- C. $y^2 - x^2 - 2xy \frac{dy}{dx} = 0$
- D. $y^2 + 2x \frac{dy}{dx} = 0$
65. The correlation coefficient lies between
- A. 0 and 1
- B. -1 and 0
- C. -1 and 1
- D. None of the above
66. The average monthly production of a factory for the first 8 months is 2500 units, the next 4 months is 1200 units. The average monthly production of the year will be
- A. 2066.55
- B. 5031.10
- C. 4021.10
- D. 3012.11
67. The equation of the curve that passes through the point (1,1) and has at every point the slope $\frac{-y}{x}$ is
- A. $xy = 1$
- B. $xy = -1$
- C. $xy = 2$
- D. $xy = -2$

68. The solution of the differential equation $\frac{d^2 y}{dx^2} + -3\frac{dy}{dx} + 2y = 0$, subject to initial conditions $y(0) = 0, y'(0) = 1$ is
- A. $e^x + e^{2x}$
 - B. $e^x - e^{2x}$
 - C. $-e^x + e^{2x}$
 - D. None of the above
69. The general solution of the differential equation $\frac{d^2 y}{dx^2} + y = x$ is
- A. $\sin x + \cos x + x$
 - B. $\sin x + \cos x + 1$
 - C. $\sin x - \cos x + x$
 - D. $\sin x - \cos x - 1$
70. The partial differential equation $u_t = ku_{xx}$ is
- A. one dimension wave equation
 - B. elliptic
 - C. one dimension heat equation
 - D. not parabolic
71. The partial differential equation $(u_{xx})^2 + u_{yy} + a(x, y)u_x + b(x, y)u - 4e^x = 0$ is
- A. linear
 - B. quasilinear
 - C. nonlinear
 - D. homogeneous

72. The solution of the IVP $\frac{dy}{dx} = x^2y - 3x^2, y(0) = 1$ is $y =$
- A. $3 + ce^{x^3/3}, c$ is a constant
 - B. $3 - 2e^{x^3/3}$
 - C. $3 + 3e^{x^3/3}$
 - D. $3 - 2e^{x^3}$
73. If $q(x) \leq 0$, then any nontrivial solution of $y'' + q(x)y = 0$
- A. can have more than one zero
 - B. can have at most one zero
 - C. cannot have any zero
 - D. has exactly one zero
74. A solution of the partial differential equation $u_t + cu_x = 0$ is $u(x, t) =$
- A. $\sin(x - t)$
 - B. $\cos(x - ct)$
 - C. $\cos(cx - t)$
 - D. $\cos xt$
75. The general solution of the partial differential equation $\frac{\partial x \partial x}{\partial x \partial y} = xy$ is
- A. $z = a \frac{x^2}{2} - \frac{y^2}{2a} + b$
 - B. $z = a \frac{x^2}{2} + \frac{y^2}{2a} - b$
 - C. $z = a \frac{x^2}{2} + \frac{y^2}{2a} + b$

D. $z = a \frac{x^2}{2} - \frac{y^2}{2a} - b$

76. The function $f : (-1, 1) \rightarrow \mathbb{R}$ defined by $f(x) = \frac{x}{1-|x|}$ is
- A. one-one but not onto
 - B. not onto
 - C. one-one and onto
 - D. neither one-one nor onto
77. For each $n \in \mathbb{N}$, let $a_n = \sum_{k=1}^n \frac{(-1)^{k-1}}{k}$. Then the sequence (a_n) is
- A. not a Cauchy sequence
 - B. a convergent sequence
 - C. not a bounded sequence
 - D. convergent to 0
78. If a function f is continuous in $[a, b]$ and differentiable (a, b) , then there exists at least one number $\theta \in (a, b)$ such that $f(a+h) = f(a) + hf'(a+\theta h)$. It is the statement of the
- A. Roll's Theorem
 - B. Lagrange's Mean Value Theorem
 - C. Cauchy's Mean Value Theorem
 - D. Taylor's Mean Value Theorem
79. The set of all polynomials with rational coefficients
- A. is not countable
 - B. is finite
 - C. does not contain $\mathbb{Q}$
 - D. countable

80. The function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = \begin{cases} x \sin\left(\frac{1}{x}\right) & \text{if } x \neq 0 \\ 0 & \text{if } x = 0 \end{cases}$.

Then at the point 0

- A. f is continuous
- B. f is not continuous
- C. f is differentiable
- D. None of the above

81. $\lim_{n \rightarrow \infty} \frac{2^{n+1} + 3^{n+1}}{2^n + 3^n}$ is equal to

- A. 3
- B. 2
- C. 1
- D. 0

82. The function $f : \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x) = \begin{cases} x-1 & \text{if } x \geq 1 \\ 1-x & \text{if } x < 1 \end{cases}$

Then at the point 1

- A. f is continuous
 - B. f is not continuous
 - C. f is differentiable
 - D. None of the above
83. The smallest number with 18 divisors is
- A. 360
 - B. 175
 - C. 185
 - D. 180

84. The value of the $\int_{|z|=2} \frac{\sin z}{\left(z - \frac{\pi}{2}\right)^2} dz$
- A. -1
B. 1
C. 0
D. π
85. $\left(\frac{1+i}{\sqrt{2}}\right)^4$ is equal to
- A. 1
B. 0
C. $\sqrt{2}$
D. -1
86. The set $\{z \in \mathbb{C} : |z - 3i| = 3\}$ geometrically represents a
- A. Parabola
B. Circle
C. Ellipse
D. Hyperbola

87. The bilinear transformation which maps the points $z_1 = 0$, $z_2 = -i$ and $z_3 = -1$ into $w_1 = i$, $w_2 = 1$ and $w_3 = 0$ respectively is

A. $\frac{-i(z+1)}{z-1}$

B. $\frac{-i(z-1)}{z+1}$

C. $\frac{-1(z+i)}{z-i}$

D. $\frac{(z+i)}{z-i}$

88. The first two terms of the Laurent's series of $f(z) = \frac{z}{(z-1)(2-z)}$, valid for $|z| > 2$, is

A. $\frac{1}{z} - \frac{3}{z^2}$

B. $-\frac{1}{z} - \frac{3}{z^2}$

C. $\frac{1}{z} + \frac{3}{z^2}$

D. None of the above

89. Z_6 is

A. a ring

B. an integral domain

C. a field

D. None of the above

90. The set of all square matrices of order 2 over the set of real number is
- A. Ring
 - B. Integral Domain
 - C. Field
 - D. None of the above
91. Let S be the set of functions $f: \mathbb{R} \rightarrow \mathbb{R}$ which are solutions to the differential equation $f''' + f' - 2f = 0$. Then S is
- A. not a vector space
 - B. a vector space of dimension greater than 3
 - C. a vector space of dimension 3
 - D. a vector space of dimension less than 3
92. The dimension of the subspace $W = \{(x_1, x_2, x_3, x_4) \in \mathbb{R}^4 : x_1 + x_2 + x_3 + x_4 = 0\}$
- A. 1
 - B. 2
 - C. 3
 - D. 4
93. Let W be the subspace spanned by $S = \{(1, 0, 0, 0), (0, 1, 0, 0), (1, 1, 0, 0), (1, 1, 1, 0), (2, 0, 3, 0)\}$. Then the dimension of W is
- A. 4
 - B. 5
 - C. 2
 - D. 3

94. The number of subsets (including the empty subset and the whole set) for a set of n elements is
- A. n
 - B. n^2
 - C. n^n
 - D. 2^n
95. Let G be the complete graph on n vertices. Then the number of edges in G is
- A. n
 - B. n^2
 - C. $2n$
 - D. $\frac{n(n-1)}{2}$
96. For $n \geq 4$, let G be a graph with n vertices and n edges. Then
- A. G is a star
 - B. G should contain a cycle
 - C. G is acyclic
 - D. G is a complete graph
97. If Z is the optimal solution of a LPP and Z' is the optimal solution of its dual, then
- A. $Z < Z'$
 - B. $Z > Z'$
 - C. $Z \neq Z'$
 - D. $Z = Z'$

98. The differential equation obtained by eliminating f from $z = f(x^2 + y^2)$ when $p = \frac{\partial f}{\partial x}$ and $q = \frac{\partial f}{\partial y}$ is
- A. $py = qx$
 - B. $pq = xy$
 - C. $px = qy$
 - D. $x = y$
99. The differential equation of the family of curves $y = e^{2x}(A \cos x + B \sin x)$ where A and B are constants is
- A. $\frac{d^2 y}{dx^2} - 4 \frac{dy}{dx} + 4y = 0$
 - B. $\frac{d^2 y}{dx^2} - 4 \frac{dy}{dx} + 5y = 0$
 - C. $\frac{d^2 y}{dx^2} - 4y \frac{dy}{dx} + 5y = 0$
 - D. $\frac{d^2 y}{dx^2} - 4 \frac{dy}{dx} + 4x = 0$
100. Let $(y - c)^2 = cx$ be the primitive of the differential equation $4x \left(\frac{dy}{dx} \right)^2 + 2x \left(\frac{dy}{dx} \right) - y = 0$. Then number of integral curve(s) which will pass through $(1, 2)$ is
- A. one
 - B. two
 - C. three
 - D. four

101. If y_1 and y_2 are two independent solutions of the differential equation $\frac{d^2 y}{dx^2} + P(x)\frac{dy}{dx} + Q(x)y = 0$, where $P(x)$ and $Q(x)$ are continuous functions of x , then which of the following is true?

A. $y_1 y_1' - y_2 y_2' = C e^{-\int P dx}$

B. $y_1 y_2' - y_2 y_1' = C e^{-\int P dx}$

C. $y_1 y_1' + y_2 y_2' = C e^{\int P dx}$

D. $y_1 y_2' + y_2 y_1' = C e^{\int P dx}$

102. Let the chances of solving a problem given to three students are $\frac{1}{2}, \frac{1}{3}, \frac{2}{5}$. The probability that the problem will be solved is

A. $\frac{1}{15}$

B. $\frac{1}{5}$

C. $\frac{3}{5}$

D. $\frac{4}{5}$

103. To measure flatness or peakedness of a distribution we use

A. skewness

B. kurtosis

C. correlation

D. standard deviation

104. A method for solving linear programming problem without using artificial variables is
- A. Two-phase method
 - B. Big-M method
 - C. Dual simplex method
 - D. Revised simplex method
105. In an assignment problem of order n , in the reduced cost matrix, the minimum number of lines needed to cover all zeros will be
- A. n
 - B. $n-1$
 - C. $n+1$
 - D. $(n-1)(n+1)$
106. The angle between two forces of equal magnitude P when their resultant also has the same magnitude P is
- A. 120°
 - B. 60°
 - C. 30°
 - D. 45°
107. The horizontal range of the projectile is maximum when the particle is projected at an angle of
- A. 90° to the horizontal
 - B. 30° to the horizontal
 - C. 60° to the vertical
 - D. 45° to the vertical

108. The period of oscillation of a simple pendulum is

A. $2\pi\sqrt{\frac{g}{l}}$

B. $\frac{\sqrt{2\pi l}}{g}$

C. $\sqrt{\frac{2\pi l}{g}}$

D. $2\pi\sqrt{\frac{l}{g}}$

109. The series $x + x^3/3! + x^5/5! + \dots$ represents the function

A. $\cos x$

B. $\cosh x$

C. $\sin x$

D. $\sinh x$

110. The value of $\frac{1}{2}(\log(1+x) - \log(1-x))$ equals

A. $\tan x$

B. $\tanh x$

C. $\tan^{-1} x$

D. $\tanh^{-1} x$

111. The value of $p(4)$, where p is the number of possible partitions of 4 is

A. 10

B. 5

C. 4

D. 1

112. Let $f(x) \in R[x] = \{a_0 + a_1x + \dots + a_nx^n : a_i \in R \text{ a ring}\}$, where n is a non-negative integer. If $f(a) = f'(a) = 0$, then $(x-a)^2$ divides
- $f(x)$
 - $f'(x)$
 - $f(x) - a$
 - $f'(x) - a$
113. Let G be a graph with $n > 2$ vertices. If all the n vertices in G form a cycle, then the degree of any vertex in G is
- 1
 - 2
 - $n-1$
 - n
114. Consider the graph G with 6 vertices given by $v_1, v_2, v_3, v_4, v_5, v_6$ and edges a, b, c, d, e, f, g, h . Then which among the following is a possibility for a path?
- $v_1 a v_2 b v_3 c v_3 d v_4 e v_2 f v_5$
 - $v_2 b v_3 d v_4 e v_2 a v_1$
 - $v_1 a v_2 b v_3 d v_4$
 - $v_2 b v_3 d v_4 h v_5 f v_2$
115. The maximum possible number of edges in a simple graph with n vertices and 2 components is
- $(n-1)(n-2)/2$
 - $n(n-1)/2$
 - $n(n+1)/2$
 - $n^2/2$

116. A Hamiltonian circuit is possessed by every graph with three or more vertices if it is
- A. connected
 - B. a tree
 - C. simple
 - D. complete
117. The rank of incidence matrix $A(G)$ of a disconnected graph G with n vertices and k components is
- A. k
 - B. $k-1$
 - C. $n-k$
 - D. $n-k+1$
118. Let T be a tree with four vertices v_1, v_2, v_3, v_4 . Then the possible number of paths between the vertices v_1 and v_4 is
- A. atleast two
 - B. atleast one
 - C. exactly one
 - D. exactly two
119. For a graph G , both the incidence matrix $A(G)$ and adjacency matrix $X(G)$ contain the entire information about G if,
- A. G has no self-loops
 - B. G has no parallel edges
 - C. G has self-loops but no parallel edges
 - D. G is simple

120. Which of the integrals does not have a definite value?

A. $\int_a^\infty \sin x \, dx, a > 0$

B. $\int_1^\infty \frac{1}{x^2} dx$

C. $\int_{-\infty}^0 e^x dx$

D. $\int_0^1 \frac{1}{\sqrt{x}} dx$

121. If p is a prime number, then $(p-1)!+1$ is

A. an odd number

B. an even number

C. a prime number

D. divisible by p

122. The curvature of a circle of radius r is

A. $\frac{1}{r^2}$

B. $-\frac{1}{r^2}$

C. r^2

D. $\frac{1}{r}$

123. The area enclosed by the curve $|x|+|y|=1$ is

A. 1

B. $\sqrt{2}$

C. 2

D. 4

124. If $(1+x)^n = C_0 + C_1x + C_2x^2 + \dots + C_nx^n$, then the value of $C_0 + 2C_1 + 3C_2 + \dots + (n+1)C_n$ is

A. $(n+2)2^{n-1}$

B. $(n+2)2^n$

C. $(n+1)2^{n-1}$

D. $(n+1)(n+2)2^n$

125. The value of $\zeta(4)$ is

A. $\pi^2/12$

B. $\pi^4/20$

C. $\pi^4/90$

D. π

126. The area under one arc of the cycloid $x = a(\theta - \sin \theta)$, $y = a(1 - \cos \theta)$ is

A. $\frac{\pi a^2}{8}$

B. $\frac{3\pi a^2}{16}$

C. $3\pi a^2$

D. $\frac{3\pi a^2}{32}$

127. The area between the parabola $y^2 = 4ax$ and the line $y = x$ is

A. $\frac{3a^2}{8}$

B. $\frac{8a^2}{3}$

C. $\frac{a^2}{8}$

D. $\frac{5a^2}{8}$

128. If $u = \tan^{-1}\left(\frac{y}{x}\right)$, then $x\frac{\partial u}{\partial x} + y\frac{\partial u}{\partial y}$ is equal to

A. $\frac{2xy}{x^2 + y^2}$

B. 1

C. 0

D. $\frac{x^2}{x^2 + y^2}$

129. The velocity of a particle moving in a straight line is given by $v^2 = se^s$, where s is the displacement in time t . The acceleration of the particle is given by

A. $(1+s)v/2$

B. $v^2/2(1+s)$

C. $v/2(s-1)$

D. $v/2(1+1/s)$

130. The directional derivative of $f(x, y) = 2x^2 + 3y^2 + z^2$ at point $(2, 1, 3)$ in the direction $\hat{i} - 2\hat{k}$ is
- A. $4/\sqrt{5}$
 B. $-4/\sqrt{5}$
 C. $\sqrt{5}/4$
 D. $-\sqrt{5}/4$
131. The value of $\iiint xyz \, dx \, dy \, dz$ taken through the positive octant of the sphere $x^2 + y^2 + z^2 = a^2$ is
- A. $\frac{a^3}{48}$
 B. $\frac{a^6}{48}$
 C. $\frac{a^6}{8}$
 D. $\frac{a^5}{48}$
132. If Σ is the boundary surface of a three dimensional region V having ζ as the unit vector along the exterior normal to the surface Σ and y is a vector function, then $\iiint_V \nabla \cdot y \, dV = \iint_{\Sigma} y \cdot \zeta \, d\Sigma$ is called
- A. Stokes theorem
 B. Gauss theorem
 C. Green's theorem
 D. Cauchy's theorem

133. The vector equation of a sphere one of whose diameters has the extremities with the position vectors a and b is
- A. $(r-a) \times (r-b) = 0$
- B. $(r+a) \times (r+b) = 0$
- C. $(r+a) \cdot (r+b) = 0$
- D. $(r-a) \cdot (r-b) = 0$
134. Consider a closed surface S surrounding a volume V . If $\vec{r}$ is the position vector of a point inside S with $\hat{n}$ the unit normal on S , the value of the integral $\iiint_V \vec{r} \cdot \hat{n} dS$ is
- A. $3V$
- B. $5V$
- C. $10V$
- D. $15V$
135. The dimension of the vector space V of all polynomials in $[x]$ of degree ≤ 20 is given by
- A. 21
- B. 20
- C. 10
- D. 8
136. If $A = (a_{ij})$ is an $n \times n$ matrix defined over a field F , then trace of A is
- A. 0
- B. I
- C. $\sum_{i=1}^n a_{ii}$
- D. $\sum_{i,j=1}^n a_{ij}$

137. If $F(x) = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix}$, then $F(x)F(y) =$

A. $F(xy)$

B. $F(x) + F(y)$

C. $F(x+y)$

D. $F(x-y)$

138. If $A = \begin{bmatrix} 1 & 0 & 0 \\ i & \frac{-1+i\sqrt{3}}{2} & 0 \\ 0 & 1+2i & \frac{-1-i\sqrt{3}}{2} \end{bmatrix}$, then the trace of A^{102} is

A. 0

B. 1

C. 2

D. 3

139. If the matrix $A = \begin{bmatrix} 3 & -1 & 1 \\ -15 & 6 & -5 \\ x & -2 & 2 \end{bmatrix}$ is singular, then the value of x is

A. 6

B. 5

C. 3

D. 2

140. If V and W are vector spaces of dimensions m and n respectively over a field F , then the dimension of the vector space of all homomorphisms of V into W is
- A. $m + n$
 - B. $m - n$
 - C. mn
 - D. m/n
141. The eigen values of a real symmetric matrix are always
- A. positive
 - B. imaginary
 - C. real
 - D. complex conjugate pairs
142. If A is an $n \times n$ matrix with diagonal entries a and other entries b , then one eigen value of A is $a - b$. Another eigen value of A is
- A. $b - a$
 - B. $nb + a - b$
 - C. $nb - a + b$
 - D. 0
143. What is the degree of the first order forward difference of a polynomial of degree n ?
- A. n
 - B. $n - 1$
 - C. $n - 2$
 - D. $n + 1$
144. The rate of convergence of bisection method is
- A. linear
 - B. faster than linear but slower than quadratic
 - C. quadratic

- D. cubic
145. The set $\mathbb{N}$ of natural numbers where $x * y = \max \{x, y\}$ is a
- A. ring
 - B. complete lattice
 - C. semi group
 - D. field
146. In a three dimensional Euclidean space, the direction cosines of a line which is equally inclined to the axes are
- A. 1, 1, 1
 - B. $1/3, 1/3, 1/3$
 - C. $1/\sqrt{3}, 1/\sqrt{3}, 1/\sqrt{3}$
 - D. $1/2, 1/2, 1/2$
147. In two dimensions, the distance between the origin and the centroid of the triangle joining the points $(-1, 0)$, $(4, 0)$ and $(0, 3)$ is
- A. 0
 - B. 1
 - C. $\sqrt{2}$
 - D. $\sqrt{3}$
148. Let P be the point $(1, 0)$ and Q a point on the locus $y^2 = 4x$. Then the locus of mid point of PQ is
- A. $y^2 + 2x + 1 = 0$
 - B. $y^2 - 2x + 1 = 0$
 - C. $x^2 - 2y + 1 = 0$
 - D. $x^2 + 2y + 1 = 0$

149. The Laplace transform of $\frac{\sin at}{at}$ is

A. $\tan\left(\frac{a}{s}\right)$

B. $\tan^{-1}\left(\frac{a}{s}\right)$

C. $\tan^{-1}\left(\frac{s}{a}\right)$

D. $\tan\left(\frac{s}{a}\right)$

150. The inverse Laplace transform of $\frac{s+2}{s^2-4s+13}$ is

A. $e^{2t} \cos 3t$

B. $\frac{4}{3}e^{2t} \sin 3t$

C. $e^{2t} \cos 3t + \frac{4}{3}e^{2t} \sin 3t$

D. $e^{2t} \cos 3t - \frac{4}{3}e^{2t} \sin 3t$
